

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS13 2717

ASyO2 | Mouse anti-human alpha-synuclein | oligomer-specific (clone number 51,24)

Product information

Immunogen synthetic peptide derived from human alpha-synuclein Glu131-Ala140

Host Mouse

Clonality Monoclonal

Subclass/isotype | IgG1

Purity Immunogen affinity purified serum in PBS pH 7.4.

Format Lyophilized

Quantity 50 μg

Reconstitution For reconstitution add 50 μl of sterile water

Storage For short time storage please and sodium azide and sro

For short time storage please add sodium azide and srote at +4°C.For long time storage store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application information

Recommended dilution 1-2 ug/ml (Dot), 2-4 ug/ml (ELISA capture), 10 ug/ml (IHC)

Expected | apparent 14 kD

1W

Confirmed reactivity | Human

Predicted reactivity | Mouse

Not reactive in No confirmed exceptions from predicted reactivity are currently known

Selected references Brännström et al. (2014). A Generic Method for Design of Oligomer-Specific Antibodies. PLoS ONE. DOI:

10.1371/journal.pone.0090857.

Application example


dot blot

Fibrils
Oligomers



Dot blot reaction of the binding capacity of ASyO2 to fibrils, monomers and oligomers. Equal amounts of each sample were spotted on a nitrocellulose membrane and then dried. The membrane was blocked with 5% non-fat milk before incubated for 1 h with anti-ASyO2 (25nM) and then with secondary antibody, anti-mouse HRP-conjugated (1:1500). The membrane was washed with PBS containing 0.25% Tween-20 before detection using ECL prime (GE Healthcare).

immunolocalization



Tissue sections from the human PD midbrain, substantia nigra, were de-waxed and rehydrated in ethanol and then incubated with ASyO2 at RT for 1h. The immunoreactivity was detected with the anti-mouse Peroxidase Reagent Kit (ImmPRESS, Vector Laboratories, Inc.) and then developed using the ImmPACT AEC Peroxidase Substrate kit (Vector Laboratories, Inc.).