

Product no **AS10 841****Donkey anti-Rabbit IgG (H&L), ALP conjugated****Product information**

Immunogen	Purified Rabbit IgG
Host	Donkey
Clonality	Polyclonal
Purity	Immunogen affinity purified donkey IgG.
Format	Liquid
Quantity	1 mg
Storage	Non-diluted antibody is stable for 4 years at 2-8°C. For storage at -20°C dilute antibody solution with an equal volume of glycerol to obtain final glycerol concentration of 50 % to prevent loss of enzymatic activity. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well but without foaming.
Additional information	ALP conjugate is supplied in 30 mM Triethanolamine, pH 7,2, 5 mM Magnesium Chloride, 0,1 mM Zinc Chloride, 1 % (w/v) BSA, Protease/IgG free, 0,05 % (w/v) of sodium azide is added as preservative

Application information

Recommended dilution	1 : 500-1 :2000 (ELISA), 1 : 50-1 : 5000 (ICC),, 1 : 20 -1 : 2000 (IHC), 1 : 500-1 :2000 (WB)
Confirmed reactivity	Rabbit IgG (H&L)
Predicted reactivity	Rabbit IgG (H&L)
Not reactive in	No confirmed exceptions from predicted reactivity are currently known
Additional information	This antibody reacts with alpha (heavy chains) on rabbit IgG and all light chains on rabbit immunoglobulins based on immunoelectrophoresis. No reactivity is observed to non-immunoglobulin rabbit serum proteins based in immunoelectrophoresis.
Selected references	Kay et al. (2014) . Elevations in Th2-initiating cytokines (IL-33, IL-25, TSLP) in lesional skin from Chronic Spontaneous ("Idiopathic") Urticaria. Br J Dermatol. 2014 Dec 18. doi: 10.1111/bjd.13621.