

**Product no** AS10 1008**Donkey anti-Rabbit IgG (H&L), HRP conjugated, min, cross-reactivity to bovine, chicken, goat, guinea pig, hamster, horse, human, mouse, rat, sheep IgG****Product information**

|                               |   |
|-------------------------------|---|
| <b>Host</b>                   | Donkey  |
| <b>Clonality</b>              | Polyclonal  |
| <b>Purity</b>                 | Immunogen affinity purified IgG.  |
| <b>Format</b>                 | Lyophilized   |
| <b>Quantity</b>               | 1 mg  |
| <b>Reconstitution</b>         | For reconstitution add 1,1 ml of sterile water, Let it stand 30 minutes at room temperature to dissolve, Prepare fresh working dilutions daily  |
| <b>Storage</b>                | Store lyophilized material at 2-8°C. For long time storage after reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1,1 ml of sterile water add 1,1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard, Be sure to mix well but without foaming. |
| <b>Additional information</b> | <p>This antibody reacts with the heavy chains on rabbit IgG and with the light chains on all rabbit immunoglobulins based on immunoelectrophoresis.</p> <p>Minimum cross-reactivity is observed to non-immunoglobulin rabbit serum proteins based on immunoelectrophoresis.</p> <p>Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.2, 10 % (w/v) BSA, Protease/IgG free and 0.1 % (v/v) Kathon CG is used as preservative. Use of sodium azide will inhibit enzymatic activity of horseradish peroxidase.</p>  |

**Application information**

|                             |   |
|-----------------------------|---|
| <b>Recommended dilution</b> | The optimal working dilution should be determined by the investigator   |
| <b>Selected references</b>  | <a href="#">Petersen</a> and Andersen (2014). Simultaneous isolation of mRNA and native protein from minute samples of cells. <i>Biotechniques</i> . 2014 May 1;56(5):229-37. doi: 10.2144/000114165. eCollection 2014. |