


Product no **AS13 2652****EF-G1 | elongation factor G1****Product information**

Immunogen	Recombinant EF-G1 protein from <i>Synechocystis</i> sp. PCC6803, P28371 , Cyanobase: slr1463
Host	Rabbit
Clonality	Polyclonal
Purity	Serum
Format	Lyophilized
Quantity	50 µl
Reconstitution	For reconstitution add 50 µl of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application information

Recommended dilution	1 : 4000 (WB)
Expected apparent MW	76,7 kDa
Confirmed reactivity	<i>Streptococcus pneumoniae</i> , <i>Synechocystis</i> sp. PCC6803
Predicted reactivity	Cyanobacteria, <i>Nannochloropsis gaditana</i> Species of your interest not listed? Contact us
Not reactive in	No confirmed exceptions from predicted reactivity are currently known
Selected references	Bárria et al. (2019). Pneumococcal RNase R globally impacts protein synthesis by regulating the amount of actively translating ribosomes. <i>RNA Biol.</i> 2019 Feb;16(2):211-219. doi: 10.1080/15476286.2018.1564616.

Application example

Left panel: from 0.15 to 5.17 µg of total protein from *Synechocystis* PCC6803 extracted with SDS-sample buffer and recombinant EF-G1 (**right panel**) were separated on 10 % SDS-PAGE and blotted 1h to **PVDF**. Blots were blocked with 5 % milk powder in TBS-T for 30 min. at room temperature (RT) with agitation. Blot was incubated in the primary antibody at a dilution of 1: 4 000 for 1h at RT with agitation. The antibody solution was decanted and the blot was rinsed briefly twice, then washed 3 times for 7 min in TBS-T at RT with agitation. Blot was incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated) diluted to 1:10 000 in for 1h at RT with agitation. The blot was washed as above and developed for 5 min with ECL according to the manufacturer's instructions. Exposure time was 2 minutes.

Courtesy of Yichen Zhang, Department of Biochemistry and Molecular Biology, University of Massachusetts, USA