


Product no **AS07 251****FtsH10 | ATP-dependent zinc metalloprotease FtsH10 (mitochondrial)****Product information**

<b>Immunogen</b>	KLH-conjugated peptide located near C-terminus chosen from sequence of <i>Arabidopsis thaliana</i> FtsH10 <a href="#">Q8VZ18</a> , <a href="#">At1g07510</a>
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Serum
<b>Format</b>	Lyophilized
<b>Quantity</b>	200 µl
<b>Reconstitution</b>	For reconstitution add 200 µl of sterile water
<b>Storage</b>	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

**Additional information** | Blue-native (2D BN/SDS-PAGE) methodology has been described in Piechota et al, 2010

**Application information**

<b>Recommended dilution</b>	1 : 1000 (WB)
<b>Expected   apparent MW</b>	84 kDa
<b>Confirmed reactivity</b>	<i>Arabidopsis thaliana</i>
<b>Predicted reactivity</b>	<i>Arabidopsis thaliana</i>
<b>Not reactive in</b>	No confirmed exceptions from predicted reactivity are currently known
<b>Selected references</b>	<p><a href="#">Kolodziejczak et al. (2018)</a>. m-AAA Complexes Are Not Crucial for the Survival of Arabidopsis Under Optimal Growth Conditions Despite Their Importance for Mitochondrial Translation. <i>Plant Cell Physiol.</i> 2018 May 1;59(5):1006-1016. doi: 10.1093/pcp/pcy041.</p> <p><a href="#">Piechota et al. (2015)</a>. Unraveling the functions of type II-prohibitins in Arabidopsis mitochondria. <i>Plant Mol Biol.</i> 2015 Apr 21.</p> <p><a href="#">Kwasniak et al. (2013)</a>. Silencing of the Nuclear RPS10 Gene Encoding Mitochondrial Ribosomal Protein Alters Translation in Arabidopsis Mitochondria. <i>Plant Cell</i>, May 30.</p> <p><a href="#">Quesada et al. (2011)</a>. Arabidopsis RUGOSA2 encodes an mTERF family member required for mitochondrion, chloroplast and leaf development. <i>Plant J.</i></p>

**Application example**

**Mitochondrial preparation** from *Arabidopsis thaliana* mitochondria was separated on 10% acrilamide gel and electrophoresis prepared according to Schägger and von Jagov (*Anl. Biochem.*, 1987, 166:368-379). After running the gel, proteins were transferred to nitrocellulose membrane using wet transfer (0.22% CAPS, pH 11). Transfer was checked by Ponceau S staining. Blot was destained by several quick washings in distilled water and 1 washing in 1X TBS (10 mM T pH 7.5, 150 mM NaCl) (10-15 min.). Blot was blocked by 1.5 hour in 5% milk in TBST (1X TBS, 0.1 20) After blocking blot was washed quickly twice in TBST and incubated 2 hours with primary antibody (dilution 1: 1000 TBST (dilution 1:1000). Washing: two quick washings in TBST and 3 x 10 min. washings in TBST. Then blot was incubated 45-60 min. with a secondary anti-rabbit antibodies conjugated to peroxidase (dilution 1:10000) in TBST. Washing: as above. After washing blot was incubated 1-2 min. in ECL solution and exposed to Kodak autoradiography film. Exposure time was 15-60 seconds.

# Agrisera

Part of Olink® Group 

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: [support@agrisera.com](mailto:support@agrisera.com)

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | [www.agrisera.com](http://www.agrisera.com)

Mitochondria were isolated as described by Urantowka et al. (Plant Mol Biol, 2005, 59:239-52). Mitochondrial pellets were suspended in 1X Laemmli buffer (5% beta-mercaptoethanol, 3.7% glycerol, 1.1% SDS, 23 mM Tris-HCl pH 6.8, 0.01% bromophenol blue), heated (95 °C, 5 min.) and centrifuged (13000 rpm, 1 min.).

Courtesy Dr. J. Piechota, University of Wrocław, Poland