

**Product no** [AS16 3356](#)**Goat anti-Human IgG Fc, F(ab)'2 Fragment, min, cross-reactivity to bovine, horse, mouse or rabbit serum proteins, mouse IgG1****Product information**

|                  |                                       |
|------------------|---------------------------------------|
| <b>Immunogen</b> | Purified Human IgG, Fc fragment       |
| <b>Host</b>      | Goat                                  |
| <b>Clonality</b> | Polyclonal                            |
| <b>Purity</b>    | Immunogen affinity purified goat IgG. |
| <b>Format</b>    | Liquid                                |
| <b>Quantity</b>  | 1,0 mg                                |
| <b>Storage</b>   | Store at 2-8 °C.                      |

|                               |   |
|-------------------------------|---|
| <b>Additional information</b> | <p>Antibody purity is &gt; 90% based on SDS-PAGE. Small amounts of intact IgG may be present. Affinity purified using solid phase Human IgG .</p> <p>Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2. 0.05% (w/v) Sodium Azide is added as a preservative.</p> <p>Based on IEP, this antibody reacts with: • heavy ( ) chains on human IgG .</p> <p>Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • light chains on all human immunoglobulins • human IgG F(ab)'2 fragment • mouse IgG1 • serum proteins from bovine, horse, mouse, or rabbit .</p> |
|-------------------------------|---|

**Application information**

|                             |   |
|-----------------------------|---|
| <b>Recommended dilution</b> | The optimal working dilution should be determined by the investigator |
| <b>Confirmed reactivity</b> | Goat anti-Human IgG Fc, F(ab)'2 Fragment                              |