

Product no **AS10 1187****Goat anti-Rat IgG (H&L), HRP conjugated****Product information****Host** | Goat**Clonality** | Polyclonal**Purity** | Immunogen affinity purified rat IgG.**Format** | Lyophilized**Quantity** | 1 mg**Reconstitution** | For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Centrifuge to remove any particulates. Prepare fresh working dilutions daily.**Storage** | Store lyophilized material at 2-8°C. For long time storage after reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1.1 ml of sterile water add 1.1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well but without foaming. Store liquid material at 2-8°C up to 6 months.**Additional information** | Concentration: 1.0 mg/ml (E 1% at 280 nm = 13.0)

Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.2, 10 % (w/v) BSA, Protease/IgG free and 0.1 % (v/v) Kathon CG as preservative.

Application information**Recommended dilution** | The optimal working dilution should be determined by the investigator**Confirmed reactivity** | Heavy chains on rat IgG and with the light chains on all rat immunoglobulins**Not reactive in** | Non-immunoglobulin rat serum proteins**Selected references** | [Saintenac et al. \(2021\)](#) A wheat cysteine-rich receptor-like kinase confers broad-spectrum resistance against *Septoria tritici blotch*. *Nat Commun.* 2021 Jan 19;12(1):433. doi: 10.1038/s41467-020-20685-0. PMID: 33469010; PMCID: PMC7815785.