

**Product no AS05 084A-FITC****PsbA | D1 protein of PSII, C-terminal, FITC conjugated****Product information**

<b>Immunogen</b>	KLH-conjugated synthetic peptide derived from available plant, algal and cyanobacterial PsbA sequences, including <i>Arabidopsis thaliana</i> UniProt: <a href="#">A4QJR4</a> , TAIR: <a href="#">AtCg00020</a> , <i>Oryza sativa</i> <a href="#">P0C434</a> , <i>Populus alba</i> <a href="#">Q14FH6</a> , <i>Physcomitrella patens</i> <a href="#">Q6YXN7</a> , <i>Chlamydomonas reinhardtii</i> <a href="#">P07753</a> , <i>Synechocystis</i> sp. <a href="#">P14660</a> and many others
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Immunogen affinity purified serum in PBS pH 7.4, conjugated to FITC.
<b>Format</b>	Liquid
<b>Quantity</b>	40 µg
<b>Storage</b>	Store at 4°C for 12-18 months. A preservative may be added for long time storage up to 2 years. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

**Additional information** | PsbA | D1 protein of PSII, C-terminal, FITC conjugated

**Application information**

<b>Expected   apparent MW</b>	38   28-30 kDa
<b>Confirmed reactivity</b>	<i>Anabaena</i> 7120, <i>Arabidopsis thaliana</i> , <i>Artemisia annua</i> , <i>Arundo</i> sp., <i>Chlamydomonas reinhardtii</i> , <i>Colobanthus quitensis</i> Kunt Bartl, <i>Craterostigma</i> sp., <i>Coscinodiscus wailesii</i> , <i>Cynara cardunculus</i> var <i>altilis</i> , <i>Ditylum brightwellii</i> , <i>Glycine max</i> , <i>Hordeum vulgare</i> , <i>Lindernia</i> sp., <i>Miscanthus x giganteus</i> , <i>Marchantia polymorpha</i> (liverwort), <i>Nicotiana benthamiana</i> , <i>Panicum miliaceum</i> , <i>Panax ginseng</i> , <i>Panicum maximum</i> , <i>Paulinella chromatophora</i> (amoeba), <i>Pinus strobus</i> , <i>Physcomitrium patens</i> , <i>Prochlorococcus</i> sp. (surface and deep water ecotype), <i>Synechococcus</i> sp. PCC 7942, <i>Spirodela polyrhiza</i> , <i>Symbiodinium</i> sp., <i>Zea mays</i>
<b>Predicted reactivity</b>	Algae (brown and red), <i>Brassica napus</i> , Conifers, Cyanobacteria, Dictos, <i>Cannabis sativa</i> , <i>Galdieria sulphuraria</i> , <i>Lactuca sativa</i> , <i>Lycopersicon esculentum</i> , <i>Medicago sativa</i> , <i>Nannochloropsis</i> sp., <i>Oryza sativa</i> , <i>Ostreococcus</i> sp. <i>Pisum sativum</i> , <i>Porphyridium purpureum</i> , <i>Sesamum indicum</i> , <i>Thalassiosira pseudonana</i> , <i>Zostera marina</i> , <i>Vitis vinifera</i> cellular [compartment marker] of thylakoid membrane
	Species of your interest not listed? <a href="#">Contact us</a>
<b>Not reactive in</b>	No confirmed exceptions from predicted reactivity are currently known
<b>Additional information</b>	The antibody is appropriate for detecting both, 24 kDa or the 10 kDa C-terminal fragments, whichever is generated under given treatment conditions. In our analysis we have seen both, ca. 24 kDa and ca. 10 kDa fragments from different samples, depending on treatments and isolation procedures.
	Rabbit anti-PsbA antibody can detect more than one band of PsbA protein, e.g. precursor and mature protein as compare to the hen anti-PsbA antibodies AS01 016.
	This antibody will detect the phosphorylated form of D1 as an alternate band to the main band on a high resolution gel.
	The antibody will bind to cross-linked proteins: D1/D2, D1/cyt b559, D1/CP43.