


Product no **AS10 704****PsbA | D1 protein of PSII, DE-loop****Product information**

<b>Immunogen</b>	KLH-conjugated synthetic peptide, amino acids 234-242 of <i>Arabidopsis thaliana</i> D1 protein UniProt: <a href="#">P83755</a> , TAIR: <a href="#">AtCg00020</a>
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Serum
<b>Format</b>	Lyophilized
<b>Quantity</b>	50 µl
<b>Reconstitution</b>	For reconstitution add 50 µl of sterile water
<b>Storage</b>	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.
<b>Additional information</b>	Antibody will detect 23 kDa N-terminal fragment

**Application information**

<b>Recommended dilution</b>	1 : 10 000, thylakoid fraction (WB)
<b>Expected   apparent MW</b>	38   28-30 kDa
<b>Confirmed reactivity</b>	<i>Arabidopsis thaliana</i> , <i>Chlamydomonas reinhardtii</i> , <i>Hordeum vulgare</i> , <i>Medicago truncatula</i> , <i>Nannochloropsis oceanica</i> strain IMET1, <i>Neochloris oleoabundans</i> UTEX 1185 (chlorophyta), <i>Nicotiana tabacum</i> , <i>Physcomitrium patens</i> , <i>Pisum sativum</i> , <i>Silene vulgaris</i> , <i>Sinapsis alba</i> , <i>Spinacia oleracea</i> , <i>Synechococcus</i> sp. PCC 7942, <i>Synechocystis</i> sp. PCC6803, <i>Triticum</i> sp.
<b>Predicted reactivity</b>	<i>Cucumis sativus</i> , <i>Glycine max</i> , <i>Nannochloropsis</i> sp., <i>Oryza sativa</i> , <i>Populus balsamifera</i> , <i>Ricinus communis</i> , <i>Zea mays</i> , <i>Vitis vinifera</i> Species of your interest not listed? <a href="#">Contact us</a>
<b>Not reactive in</b>	No confirmed exceptions from predicted reactivity are currently known
<b>Additional information</b>	Antibody is recognizing a 23 kDa fragment in spinach and Arabidopsis thylakoids for usage on total cell extracts the dilution needs to be determined experimentally
<b>Selected references</b>	<a href="#">Lu</a> et al. (2021). Role of an ancient light-harvesting protein of PSI in light absorption and photoprotection. <i>Nat Commun.</i> 2021 Jan 29;12(1):679. doi: 10.1038/s41467-021-20967-1. PMID: 33514722; PMCID: PMC7846763. (blue-native PAGE) <a href="#">Mazur</a> et al. (2021) The SnRK2.10 kinase mitigates the adverse effects of salinity by protecting photosynthetic machinery. <i>Plant Physiol.</i> 2021 Dec 4;187(4):2785-2802. doi: 10.1093/plphys/kiab438. PMID: 34632500; PMCID: PMC8644180. <a href="#">Rantala</a> et al. (2020). PGR5 and NDH-1 systems do not function as protective electron acceptors but mitigate the consequences of PSI inhibition. <i>Biochim Biophys Acta Bioenerg.</i> 2020 Jan 11;1861(3):148154. doi: 10.1016/j.bbabi.2020.148154. <a href="#">Grieco</a> et al. (2020). Adjustment of photosynthetic activity to drought and fluctuating light in wheat. <i>Plant Cell Environ.</i> 2020 Mar 16. doi: 10.1111/pce.13756. <a href="#">Rantala</a> and Tikkanen et al. (2018). Phosphorylation-induced lateral rearrangements of thylakoid protein complexes upon light acclimation. <i>Plant Direct</i> Vol. 2, Issue 2.

**Application example**



Thylakoid membranes from Arabidopsis (0.05-0.75 µg of Chl) were separated on 14%AA+ 6M urea gels and blotted 1 h to PVDF. Blots were blocked immediately following transfer in 5% milk solution for 1 h at room temperature with agitation. Blots were incubated in the primary antibody at a dilution of 1:20 000 o.n. at 4°C with agitation. The antibody solution was decanted and the blot was rinsed briefly twice, then washed once for 15 min and 3 times for 5 min in TBS-T at room temperature with agitation. Blots were incubated in secondary antibody, HRP conjugated, diluted to 1:5 000. The blots were washed as above and developed for 5 min with ECL-Plus detection reagent according to the manufacturers instructions. Exposure time was 1 min in CCD camera Fuji4000.

Courtesy Professor Cornelia Spetea-Wiklund, University of Ghotenburg, Sweden