

Product no **AS09 527B****AGO1 | Argonaute 1 (40 µg, Biotin conjugated)****Product information****Immunogen** | KLH-conjugated N-terminal peptide of *Arabidopsis thaliana* AGO1 UniProt: [O04379](#), TAIR: [At1g48410](#).**Host** | Rabbit**Clonality** | Polyclonal**Purity** | Immunogen affinity purified serum in PBS pH 7.4, conjugated to biotin.**Format** | Liquid**Quantity** | 40 µg**Storage** | Store at 4°C for 12-18 months. A preservative may be added for long time storage up to 2 years.**Additional information** | Antibody binds microRNA and tasiRNAs, preference for 21nt miRNAs with 5'U.[TCA acetone total protein precipitation method](#)**Application information****Recommended dilution** | 1 : 1000 (WB)**Expected | apparent MW** | 116.4 | 130 kDa**Confirmed reactivity** | *Arabidopsis thaliana***Predicted reactivity** | *Brassica pekinensis*, *Capsella rubella*, *Malus domestica*, *Pisum sativum*, *Ricinus communis*, *Vitis vinifera*Species of your interest not listed? [Contact us](#)**Not reactive in** | *Chlamydomonas reinhardtii***Additional information** | AGO expression may be tissue specific and using floral tissue is recommended where most of the AGOs are expressed the highest. Use of proteasome inhibitors as MG132 can help to stabilize AGO proteins during extraction procedure.

The AGO1 antibody is extremely specific to AGO1 and does not cross-react with other antibodies. The evidence is 1) the peptide to which it was raised is at the very N-terminus of the protein and is not present in other AGOs 2) aAGO1 does not cross react with the AGOs which are overexpressed (AGO2, AGO3, AGO4, AGO5, AGO6, AGO9) using a western blot.