

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS16 4057

ClpR4 | ATP-dependent Clp protease proteolytic subunit-related protein 4 (chloroplastic)

Product information

Immunogen BSA-conjugated peptide derived from ClpR4 of Arabidopsis thaliana, TAIR: AT4G17040, UniProt: Q8LB10

Host Rabbit

Clonality Polyclonal

Purity Serum

Format Lyophilized

Quantity 50 ul

Reconstitution For reconstitution add 50 μl of sterile water

Storage Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to

the cap or sides of the tube.

Application information

Recommended dilution 1:4000 (WB)

Expected | apparent 33,4 | 24,5 kDa

MW

Confirmed reactivity Arabidopsis thaliana

Predicted reactivity Brassica oleracea, Cajanus cajan, Capsella rubella, Coffea canephora, Cucumis sativus, Erythranthe guttata, Eutrema

salsugineum, Genlisea aurea, Glycine max, Jatropha curcas, Medicago truncatula, Morus notabilis, Populus

trichocarpa, Ricinus communis, Solanum lycopersicum, Solanum tuberosum, Vitis vinifera

Species of your interest not listed? Contact us

Not reactive in Zea mays

Additional information For western blot detection image refer to the article below

Selected references

Siögren et al. (2004). Inactivation of the clpC1 gene encoding a chloroplast Hsp100 molecular chaperone causes growth retardation, leaf chlorosis, lower photosynthetic activity, and a specific reduction in photosystem content. Plant

Physiol. 2004 Dec;136(4):4114-26. Epub 2004 Nov 24.