

Product no **AS09 414**

Cytokinin | Trans-zeatin riboside (1mg)

Product information

Immunogen	BSA-conjugated, via ribose, trans-zeatin riboside
Host	Rabbit
Clonality	Polyclonal
Purity	Total IgG. Protein G purified in PBS.
Format	Lyophilized
Quantity	1 mg
Reconstitution	For reconstitution add 70 µl of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.
Additional information	Cytokinin Trans-zeatin riboside (1mg)

Application information

Recommended dilution	2-10 µl/15 ml; Specific information about dilution is going to be included on the vial
Confirmed reactivity	Trans-zeatin riboside
Predicted reactivity	Trans-zeatin riboside
Not reactive in	No confirmed exceptions from predicted reactivity are currently known
Additional information	For ELISA kit - please inquire
Selected references	Ferreira et al. (2019). Enzyme-mediated metabolism in nutritive tissues of galls induced by <i>Ditylenchus gallaeformans</i> (Nematoda: Anguinidae). <i>Plant Biol (Stuttg)</i> . 2019 May 18. doi: 10.1111/plb.13009.