


Product no **AS07 205****FtsH4 | ATP-dependent zinc metalloprotease FtsH4 (mitochondrial)****Product information**

<b>Immunogen</b>	KLH-conjugated peptide derived from sequence of <i>Arabidopsis thaliana</i> FtsH4 UniProt: <a href="#">O80983</a> , TAIR: <a href="#">At2g26140</a>
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Immunogen affinity purified serum in PBS pH 7.4.
<b>Format</b>	Lyophilized
<b>Quantity</b>	200 µg
<b>Reconstitution</b>	For reconstitution add 100 µl of sterile water
<b>Storage</b>	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

**Application information**

<b>Recommended dilution</b>	1 : 500 (WB)
<b>Expected   apparent MW</b>	77   72 kDa
<b>Confirmed reactivity</b>	<i>Arabidopsis thaliana</i> , <i>Brassica oleracea</i> var. botrytis
<b>Predicted reactivity</b>	<i>Brassica</i> sp. Species of your interest not listed? <a href="#">Contact us</a>
<b>Not reactive in</b>	<i>Solanum lycopersicum</i>
<b>Selected references</b>	<a href="#">Opalińska et al. (2017)</a> . Identification of Physiological Substrates and Binding Partners of the Plant Mitochondrial Protease FTSH4 by the Trapping Approach. <i>Int J Mol Sci.</i> 2017 Nov 18;18(11). pii: E2455. doi: 10.3390/ijms18112455. <a href="#">Dolzbłasz et al. (2016)</a> . The mitochondrial protease AtFTSH4 safeguards Arabidopsis shoot apical meristem function. <i>Sci Rep.</i> 2016 Jun 20;6:28315. doi: 10.1038/srep28315. <a href="#">Rurek et al. (2015)</a> . Biogenesis of mitochondria in cauliflower ( <i>Brassica oleracea</i> var. botrytis) curds subjected to temperature stress and recovery involves regulation of the complexome, respiratory chain activity, organellar translation and ultrastructure. <i>Biochim Biophys Acta.</i> 2015 Jan 21. pii: S0005-2728(15)00016-X. doi: 10.1016/j.bbabi.2015.01.005. <a href="#">Zhang et al. (2014)</a> . Perturbation of auxin homeostasis caused by mitochondrial FtSH4 gene-mediated peroxidase accumulation regulates Arabidopsis architecture. <i>Mol Plant.</i> 2014 Jan 30. <a href="#">Kwasniak et al. (2013)</a> . Silencing of the Nuclear RPS10 Gene Encoding Mitochondrial Ribosomal Protein Alters Translation in Arabidopsis Mitochondria. <i>Plant Cell</i> , May 30.

**Application example**

**Total protein** from *Arabidopsis thaliana* mitochondria (20 µg) were separated on 10% acrilamide gel and electrophoresis prepared according to Schägger and von Jagov (Anl. Biochem., 1987, 166:368-379). After running the gel, proteins were transferred to nitrocellulose membrane using wet transfer (0.22% CAPS, pH 11). Transfer was checked by Ponceau S staining. Blot was destained by several quick washings in distilled water and 1 washing in 1X TBS (10 mM T pH 7.5, 150 mM NaCl) (10-15 min.). Blot was blocked by 1.5 hour in 5% milk in TBST (1X TBS, 0,1 20) After blocking blot was washed quickly twice in TBST and incubated 2 hours with primary antibody (dilution 1: 1000 TBST (dilution 1:1000). Washing: two quick washings in TBST and 3 x 10 min. washings in TBST. Then blot was incubated 45-60 min. with a secondary anti-rabbit antibodies conjugated to peroxidase (dilution 1:10000) in TBST. Washing: as above. After washing blot was incubated 1-2 min. in chemiluminescent solution and exposed to Kodak autoradiography film. Exposure time was 15-60 seconds.

Mitochondria were isolated as described by Urantowka et al. (Plant Mol Biol, 2005, 59:239-52). Mitochondrial pellets were suspended in 1X Laemmli buffer (5% beta-mercaptoethanol, 3.7% glycerol, 1.1% SDS, 23 mM Tris-HCl pH 6.8, 0.01% bromophenol blue), heated (95°C, 5 min.) and centrifuged (13000rpm, 1 min.).

Courtesy Dr. J. Piechota, University of Wrocław, Poland