

Product no **AS16 3270****Goat anti-Dog IgG (H&L), ALP conjugated****Product information**

| | |
|------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Immunogen | Purified Dog IgG, whole molecule, |
| Host | Goat |
| Clonality | Polyclonal |
| Purity | Immunogen affinity purified goat IgG. |
| Format | Liquid |
| Quantity | 0,5 mg |
| Storage | Store non-diluted antibody at 2-8°C. For storage at -20 °C, dilute with an equal volume of glycerol to prevent loss of enzymatic activity. Prepare working dilutions prior to use and then discard. |

| | |
|-------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Additional information | Antibody purity is > 95% based on SDS-PAGE. Affinity purified using solid phase Dog IgG (H&L). Antibody is supplied in 30 mM Triethanolamine, pH 7.2, 5 mM Magnesium Chloride, 0.1 mM Zinc Chloride, 1 % (w/v) BSA, Protease/IgG free. 0.05% (w/v) Sodium Azide is added as a preservative. Based on IEP, this antibody reacts with: • heavy () chains on dog IgG • light chains on all dog immunoglobulins . Based on IEP, no reactivity is observed to: • non-immunoglobulin dog serum immunoglobulins . |
|-------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

Application information

| | |
|-----------------------------|----------------------------------|
| Recommended dilution | 1 : 20-1 : 2000 (Flow cyt), (IL) |
| Confirmed reactivity | Dog IgG |