

Product no **AS10 1221**

Goat anti-Guinea pig IgG (H&L), TRITC conjugated

Product information

Host | Goat**Clonality** | Polyclonal**Purity** | Immunogen affinity purified IgG.**Format** | Lyophilized**Quantity** | 1 mg**Reconstitution** | For reconstitution add 1,1 ml of sterile water, Let it stand 30 minutes at room temperature to dissolve, Prepare fresh working dilutions daily**Storage** | Store lyophilized material at 2-8°C. For long time storage after reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1,1 ml of sterile water add 1,1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well but without foaming.**Additional information** | This antibody reacts with the heavy chains on guinea pig IgG and with the light chains on all guinea pig immunoglobulins based on immunoelectrophoresis.

Minimum cross-reactivity is observed to non-immunoglobulin guinea pig serum proteins based on immunoelectrophoresis.

Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.2, 1% (w/v) BSA, protease/IgG free and 0.05 % (w/v) sodium azide as preservative.

Application information

Recommended dilution | The optimal working dilution should be determined by the investigator,