

Product no **AS16 3468****Goat anti-Mouse IgG (H&L), F(ab)'2 fragment, TRITC conjugated, min, cross-reactivity to bovine, goat, human, rabbit, or rat IgG****Product information****Immunogen** | Purified Mouse IgG, whole molecule**Host** | Goat**Clonality** | Polyclonal**Purity** | Immunogen affinity purified goat IgG.**Format** | Lyophilized**Quantity** | 0,5 mg**Reconstitution** | For reconstitution add 0,55 ml of sterile water, Let it stand 30 minutes at room temperature to dissolve, Prepare fresh working dilutions daily**Storage** | Store at 2-8°C.

Additional information | Antibody purity is > 90% based on SDS-PAGE Small amounts of intact IgG may be present.
 Affinity purified using solid phase Mouse IgG .
 Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free.

Based on IEP, this antibody reacts with: • heavy () chains on mouse IgG • light chains on all mouse immunoglobulins .
 Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • bovine, goat, human, rabbit or rat IgG .

Application information**Recommended dilution** | 1 : 20-1 : 2000 (Flow cyt), (IL)**Confirmed reactivity** | Mouse IgG (H&L), F(ab)'2 Fragment