

## Product no **AS15 3050A** RD | N-terminal arginylation


### Product information

<b>Immunogen</b>	KLH-conjugated synthetic peptide: H-RDHKHKANQHMSVC-NH <sub>2</sub>
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Immunogen affinity purified serum in PBS pH 7.4.
<b>Format</b>	Lyophilized
<b>Quantity</b>	2 x 25 µg
<b>Reconstitution</b>	For reconstitution add 25 µl of sterile water to each tube
<b>Storage</b>	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.
<b>Additional information</b>	Antibodies are purified using subtractive purification method.  MG132 or epoxomycin are recommended to use to inhibit proteasome and significantly increase signal from the arginylated proteins.  For exact protocol of dot blot and SPOT assay, please <a href="#">inquire</a> .

### Application information


**Recommended dilution** | 1 : 1000 (Dot)

#### Application information



- Membrane: PVDF, Hybond 0.45 µm (GE Healthcare)
- 0.5 µg synthetic peptide per spot
- Membrane was blocked overnight in TBST, 4% Blocking Agent (GE Healthcare)
- Primary antibodies: 1:10 000 in TBST 2% Blocking Agent
- Secondary antibody (AS09 602): 1:25 000 in TBST 2% Blocking Agent
- Detection: ECL


Courtesy of Dr. Sebastian Hoernstein, Faculty of Biology, University of Freiburg; Germany



Dotblot of peptide variants probed with anti-RD antibodies<sup>1</sup>. The respective volumes of a 0.1 mM aqueous peptide solution were mixed with each

1  $\mu$ l of 0.1% SDS in 0.5 M Na-PO<sub>4</sub> (according to Canas et al. 1993, Anal Biochem. 211), dried in a vacuum concentrator and reconstituted in 1  $\mu$ l of H<sub>2</sub>O. The peptide solutions were spotted onto a PVDF membrane and air-dried overnight. The dried membrane was subjected to semi-dry western blot electrotransfer (0.85 mA/cm<sup>2</sup>, 3 min), blocked with 4% skim milk powder in TBST overnight and probed with anti-RD antibody (at 0.4  $\mu$ g/ml in TBST) and subsequently with HRP coupled anti-sheep antibody. Detection with ECL substrate of femtogram sensitivity, 3 min exposure time.

Courtesy Dr. Nico Dissmeyer, Leibniz Institute of Plant Biochemistry (IPB), Germany



SPOT assay with anti-RD antibody. 11- to 17mer peptides of the indicated sequences were synthesized on a PEG-derivatized cellulose membrane. The SPOT membrane was blocked overnight in TBST (50 mM TRIS-HCl, pH = 7.9; 0.15 M NaCl; 0.1% Tween-20) containing 7.5% skim milk powder. The next day, purified anti-Arg-RD21 was incubated with the membrane in TBST at 0.4  $\mu$ g/ml for 1 hour under agitation at room temperature. Subsequently, the membrane was washed 3x 10min in TBST and electrotransferred to a PVDF membrane (30min, 0.54 mA per cm<sup>2</sup>; blotting system according to Kyhse-Andersen 1984, J Biochem Biophys Methods 10(3-4)). Immobilized antibody was probed with HRP coupled anti-sheep secondary antibody and detected with ECL substrate of picogram sensitivity and 3 min exposure time. Letters in the grid represent N-terminal amino acids in 1st and 2nd position; grey background: variants of the peptide antigen used by Wong et al. 2007, PLoS Biol. 5(10).

Courtesy Dr. Nico Dissmeyer, Leibniz Institute of Plant Biochemistry (IPB), Germany