

Product no **AS19 4262****RPT2a | Regulatory particle triple-A ATPase subunit 2a****Product information**

Immunogen	Recombinant, full length RPT2a of <i>Arabidopsis thaliana</i> RPT2a protein sequence UniProt: Q9SZD4-1 , TAIR: At4g29040 overexpressed in <i>E.coli</i> , purified from a gel piece
Host	Rabbit
Clonality	Polyclonal
Purity	Serum
Format	Lyophilized
Quantity	50 µl
Reconstitution	For reconstitution add 50 µl, of sterile water
Storage	Store lyophilized/reconstituted at -20°C (short term, months) or at -80°C (long term, years) ; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application information

Recommended dilution	1 : 3000 (WB)
Expected apparent MW	49,4 50 kDa
Confirmed reactivity	<i>Arabidopsis thaliana</i>
Predicted reactivity	<i>Brassica napus</i> , <i>Cajanus cajan</i> , <i>Gossypium hirsutum</i> , <i>Mucuna pruriens</i> , <i>Noccaea caerulea</i> , <i>Vigna radiata</i> var. <i>radiata</i> Species of your interest not listed? Contact us
Not reactive in	No confirmed exceptions from predicted reactivity are currently known
Additional information	This antibody is also recognizing RPT2b protein. Recommended western blot conditions: SDS-PAGE, transfer to nitrocellulose, blocking 10% non-fat milk. Diluent for both primary and secondary antibodies PBS containing 0.2% Tween 20 and 1% BSA. For an image of western blot detection, refer to: Smalle et al. (2002).
Selected references	Meng , Wang, Hao, et al. (2023) RNA-binding protein MAC5A interacts with the 26S proteasome to regulate DNA damage response in Arabidopsis. <i>Plant Physiol.</i> 2023;191(1):446-462. doi:10.1093/plphys/kiac511 Smalle et al. (2002). Cytokinin growth responses in Arabidopsis involve the 26S proteasome subunit RPN12. <i>Plant Cell</i> 14, 17-32.